

**Erratum: Lindemann criterion and vortex lattice phase transitions in type-II superconductors
[Phys. Rev. B **69**, 024501 (2004)]**

Jan Kierfeld and Valerii Vinokur

(Received 26 October 2004; published 6 January 2005)

DOI: 10.1103/PhysRevB.71.029901

PACS number(s): 74.25.Qt, 74.25.Sv, 99.10.Cd

We want to point out that careful inspection and comparison of the results of our work for the locus of the amorphization transition lines for NbSe and those of Ref. 1 shows that both results practically coincide (up to an unessential correction factor), contrary to what was stated in our paper. The seeming discrepancy stemmed from the definition of the characteristic length $L_0 = \epsilon a (1-b)^{-1/2}$ [for $a < \lambda_{ab}$, see Eq. (10)] used in Ref. 1, which, unlike our definition, did not include the high-field correction factor $(1-b)^{-1/2}$. However, a similar factor had been inserted at a later stage of the calculation of Ref. 1 [cf. eq. (8) of Ref. 1], such that the final results for the phase boundaries become practically identical to our corresponding expressions. Therefore, we wish to withdraw our criticism of the results of Ref. 1 for the amorphization transition lines as unjustified. The results of our paper remain unaffected.

The authors would like to thank G. P. Mikitik and E. H. Brandt for sharing their notes on the comparison with their results.

¹G. P. Mikitik and E. H. Brandt, Phys. Rev. B **64**, 184514 (2001).